



BUTTOCK AUGMENTATION



Dr. Rajat Gupta

Board Certified Plastic
Surgeon, RG Aesthetics, India

What Is Buttock augmentation?

Buttock augmentation, medically known as gluteal augmentation, is used to improve the contour, size and/or shape of the buttocks. This can be done using implants or fat grafting technique. In some cases, the surgeon might do a combination of the two.

Buttock Augmentation procedure is done if:

- You want to Increase fullness on your buttocks
- You desire roundness and lifted projection of your buttocks
- You want to Improve the balance of your figure
- You want to enhance your self-image and self-confidence
- You have asymmetrical or flat buttocks

What Buttock Augmentation Can / Can't Do

✓ Can

- Increase fullness and roundness of your butt or buttocks
- Make your buttocks project more and stick out more
- Subsequently, improve the balance of your figure
- Enhance your silhouette into a more hourglass-like shape
- Enhance your self-image and self-confidence

✓ Can't

- Completely change the natural contour of your body
- Increase the height of your buttocks unnaturally
- Increase their size exponentially
- Change your entire silhouette


What is the Procedure of Buttock Augmentation?

Anaesthesia:

- Local or General
-

Procedure:

- Excess fat is suctioned out from other body parts using liposuction techniques.



- EVL (expansion vibration lipofilling) technique is applied to contour & shape the buttocks.
- 4- to 5-mm angled basket cannulas are used and attached to a power-assisted liposuction unit.
- Fat is deposited into the expanded area with the basket cannula
- Depending on the body size and gluteal dimensions of the patient, injection volumes range from 450 to 1100 mL or more per side.

Recovery After Buttock Augmentation?

- Outpatient procedure
- You may have some swelling, bruising & discomfort in the first 24-48 hours. This is only temporary and diminishes as the recovery proceeds.
- The initial recovery time is 1-3 days. Any swelling or soreness subsides within the first few weeks.
- You can resume work within a week and exercise within a month or as per your doctor's advice.
- Avoid sudden movements, lifting weights, etc., for approximately two week.
- Use compression garment for minimum 6 weeks after the surgery for the breasts to maintain the contours achieved immediately after the surgery.
- By one month, bruising, discomfort and most swelling will disappear and you will feel "back to normal".



Liposuction, Waist Contouring and Brazilian Buttock Lift



Liposuction and Fat Grafting to Buttocks



Butt Fat Grafting



Fat Grafting Buttocks



Fat Grafting Buttock





About Dr. Rajat Gupta



Dr. Rajat Gupta is a gold-medallist, board-certified plastic surgeon in India. As the National Secretary of the Indian Association of Aesthetic Plastic Surgery (IAAPS), he has been accredited with consistently raising the standards of aesthetic surgery in the country.

He also consistently trains prospective plastic surgeons coming from all over the world to learn the finesse with which he performs every procedure, using refined cosmetic surgery techniques and technologies.

His clientele includes the who's who of the Indian film industry, alongside prominent politicians and several international patients.

Today, Dr. Gupta sports unparalleled expertise in cosmetic surgery with a keen eye for detail and an in-depth understanding of aesthetics.

To book an appointment,
call: +91-9251711711 | email: contact@dr Rajat Gupta.com